



ELSEVIER

Carbohydrate Research 296 (1996) C13-C14

CARBOHYDRATE
RESEARCH

Contents

| | |
|--|-----|
| Complete ^1H and ^{13}C resonance assignments of a 21-amino acid glycopeptide prepared from human serum transferrin J. Lu and H. van Halbeek (Athens, GA, USA) | 1 |
| Cyclolaminarinose. A new biologically active β -(1 → 3) cyclic glucan P.E. Pfeffer, S.F. Osman, A. Hotchkiss (Wyndmoor, PA, USA), A.A. Bhagwat (College Park, MD, USA), D.L. Keister (Beltsville, MD, USA) and K.M. Valentine (Philadelphia, PA, USA) | 23 |
| Conformational studies of <i>myo</i> -inositol phosphates L.G. Barrientos and P.P.N. Murthy (Houghton, MI, USA) | 39 |
| Structural studies of the major glycolipid from <i>Saccharopolyspora</i> genus A. Gamian, H. Mordarska (Wroclaw, Poland), I. Ekiel (Montreal, Canada), J. Ulrich (Grenoble, France), B. Szponar (Wroclaw, Poland) and J. Defaye (Grenoble, France) | 55 |
| Crystal structures of hexakis-(2,6-di- <i>O</i> -methyl)-cyclomaltohexaose (dimethyl- α -cyclodextrin) crystallized from acetone, and crystallized from hot water T. Steiner (Berlin, Germany), F. Hirayama (Kumamoto, Japan) and W. Saenger (Berlin, Germany) | 69 |
| Full ^1H NMR assignment and detailed <i>O</i> -acetylation patterns of capsular polysaccharides from <i>Neisseria meningitidis</i> used in vaccine production X. Lemercier and C. Jones (South Mimms, UK) | 83 |
| The preparation of racemic and enantiomerically pure <i>myo</i> -inositol derivatives as intermediates for the synthesis of phosphatidylinositol 3-, 3,4-bis-, and 3,4,5-tris-phosphates and for the synthesis of analogues of 1 <i>D</i> - <i>myo</i> -inositol 1,3,4,5-tetrakisphosphate T. Desai, J. Gigg, R. Gigg and E. Martín-Zamora (London, UK) | 97 |
| <i>N</i> -Trichloroethoxycarbonyl-glucosamine derivatives as glycosyl donors W. Dullenkopf, J.C. Castro-Palomino, L. Manzoni and R.R. Schmidt (Konstanz, Germany) | 135 |
| Synthesis of α - and β -linked tyvelose epitopes of the <i>Trichinella spiralis</i> glycan: 2-Acetamido-2-deoxy-3- <i>O</i> -(3,6-dideoxy- <i>D</i> -arabino-hexopyranosyl)- β - <i>D</i> -galactopyranosides M.A. Probert, J. Zhang and D.R. Bundle (Edmonton, Canada) | 149 |
| Synthesis of a hexasaccharide acceptor corresponding to the reducing terminus of mycobacterial 3- <i>O</i> -methylmannose polysaccharide (MMP) W. Liao and D. Lu (Beijing, PR China) | 171 |
| The structure of stewartan, a capsular polysaccharide produced by <i>Erwinia stewartii</i> strain DC283 B.Y. Yang, J.S.S. Gray and R. Montgomery (Iowa City, IA, USA) | 183 |

| | |
|--|-----|
| Synthesis of α -D-GlcP-(1 \rightarrow 2)- α -D-GlcP-(1 \rightarrow 3)- α -D-GlcP-O-(CH ₂) ₈ COOCH ₃ for use in the assay of α -glucosidase I activity C.H. Scaman, O. Hindsgaul, M.M. Palcic and O.P. Srivastava (Edmonton, Canada) | 203 |
| Mechanism of anomeration of cyclohexyl 2-deoxy-3,4,6-tri-O-methyl-2-(N-methylacetamido)- α - and β -D-hexopyranosides under reductive-cleavage conditions Y.M. Ahn and G.R. Gray (Minneapolis, MN, USA) | 215 |
| Synthesis of tetrasaccharides as possible metastatic inhibitors X.X. Zhu, P.Y. Ding and M.S. Cai (Beijing, PR China) | 229 |
| Sulfated galactans from Australian specimens of the red alga <i>Phaeocarpus peperocarpus</i> (Gigartinales, Rhodophyta) M.-L. Liao, A. Chiovitti, S.L.A. Munro (Parkville, Australia), D.J. Craik (St. Lucia, Australia), G.T. Kraft and A. Bacic (Parkville, Australia) | 237 |
| Structural features of the pectic polysaccharides isolated from retted hemp bast fibres M.R. Vignon and C. Garcia-Jaldon (Grenoble, France) | 249 |
| Enzymatic properties of α -D-galactosidase from <i>Trichoderma reesei</i> A.N. Savel'ev, F.M. Ibatyllin, E.V. Eneyeskaya, A.M. Kachurin and K.N. Neustroev (St. Petersburg, Russia) | 261 |
| <i>Notes</i> | |
| An improved method for the preparation of some ethyl 1-thioglycosides S.K. Das and N. Roy (Calcutta, India) | 275 |
| Synthesis and semisynthesis of α -D-mannopyranosyl-(1 \rightarrow 2)- α -D-mannopyranose octaacetate Z. Szurmai and L. Jánossy (Debrecen, Hungary) | 279 |
| In vivo conversion of 6-O-sulfo-L-galactopyranosyl residues into 3,6-anhydro-L-galactopyranosyl residues in <i>Gracilaria chilensis</i> Bird, McLachlan <i>et al.</i> Oliveira J.A. Hemmingson, R.H. Furneaux and H. Wong (Lower Hutt, New Zealand) | 285 |
| <i>Corrigendum</i> | |
| Crystal structure of an adduct of sarcosine with sucrose [Carbohydrate Research 287 (1996) 117–122] R.V. Krishnakumar and S. Natarajan (Madurai, India) | C1 |
| <i>Book review</i> | C3 |
| <i>Author index</i> | C7 |
| <i>Subject index</i> | C9 |
| <i>Contents</i> | C13 |